



King's Research Portal

DOI:

[10.1016/j.schres.2010.07.019](https://doi.org/10.1016/j.schres.2010.07.019)

Document Version

Early version, also known as pre-print

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Kempton, M. (2010). Response to letter by Scott Kaplan. *Schizophrenia Research*, 124(1-3), 245.
<https://doi.org/10.1016/j.schres.2010.07.019>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

We thank Scott Kaplan for his interest in our study and his comments. There are two key questions to consider, 1) Is it valid to compare the effect sizes from a meta-analyses of first episode patients to a meta-analysis of chronic patients with schizophrenia? 2) If this is a valid comparison, would one expect to see a significant difference in the effect sizes if progressive changes exist?

1) Vita et al, (2006) compared the results of their meta-analysis of first episode patients to meta-analyses of mainly chronic patients with schizophrenia (Wright, et al. 2000) and conclude that the effect sizes were similar. However comparing the meta-analyses of Vita et al. to Wright et al. is problematic because of the following reasons:

- a) The statistical models used to calculate the pooled effect size were different as Wright et al. used a random effects meta-analysis and Vita et al. used a fixed effects meta-analysis.
- b) Wright et al. included first episode studies in addition to studies of chronic patients.

To circumvent these problems we have conducted a new meta-analysis of studies measuring the lateral ventricles included in Wright et al, Vita et al and in an additional first episode meta-analysis by Steen et al (2006). We used the same methodology as our previous cross-sectional meta-analysis (Kempton, et al. 2008) and performed a stratified meta-analysis directly comparing the effect size from chronic (n=21 studies, mean duration of illness=12 years) and first episode studies (n=11). There was no significant difference between the effect sizes ($p=0.54$).

2) We argue that a significant difference would not be expected in the above comparison due to the large increase in inter-study and inter-subject variance associated with meta-analyses of cross-sectional studies. Longitudinal studies remove variance in baseline ventricular volumes and increase statistical power to detect changes over time. Thus a meta-analysis of longitudinal changes is much more sensitive to progressive changes than comparing effect sizes from cross-sectional studies. In conclusion although the comparison between Wright et al and Vita et al is not significant, we can be confident that ventricular dilation is present both in first episode and chronic patients, and that progressive ventricular enlargement occurs in schizophrenia.

References

- Kempton, M. J., Geddes, J. R., Ettinger, U., Williams, S. C., Grasby, P. M., 2008. Meta-analysis, database, and meta-regression of 98 structural imaging studies in bipolar disorder. *Arch Gen Psychiatry* 65 (9), 1017-32.
- Steen, R. G., Mull, C., McClure, R., Hamer, R. M., Lieberman, J. A., 2006. Brain volume in first-episode schizophrenia: systematic review and meta-analysis of magnetic resonance imaging studies. *Br J Psychiatry* 188 510-8.
- Vita, A., De Peri, L., Silenzi, C., Dieci, M., 2006. Brain morphology in first-episode schizophrenia: a meta-analysis of quantitative magnetic resonance imaging studies. *Schizophr Res* 82 (1), 75-88.
- Wright, I. C., Rabe-Hesketh, S., Woodruff, P. W., David, A. S., Murray, R. M., Bullmore, E. T., 2000. Meta-analysis of regional brain volumes in schizophrenia. *Am J Psychiatry* 157 (1), 16-25.

Matthew J Kempton
King's College London
Centre for Neuroimaging Sciences
Institute of Psychiatry
UK